# Pediatric COVID-19 Vaccines

Kruti Acharya M.D.

## Agenda

### COVID-19 – Where We Have Been and Where We Are Now

### COVID-19 Vaccine Schedule

Resources

## COVID-19

Children 6 months – 4 years Over 2 million cases Over 20, 000 hospitalizations Over 200 deaths Children 5 years – 17 years Over 10 million cases Over 45,000 hospitalizations Over 600 deaths

3



# Where We Are Now

### Approval by Age Group



## Status of COVID-19 Vaccinations in Children

### One Dose

Fully Vaccinated

1,131,814	Number of 5–17-year- olds in Illinois with one dose	1,035,474	Number of 5–17-year- olds in Illinois fully vaccinated
41.23%	Percentage of 5–11- year-olds in Illinois with one dose	37.32%	Percentage of 5–11- year-olds in Illinois fully vaccinated
68.15%	Percentage of 12–17- year-olds in Illinois with one dose	62.80%	Percentage of 12–17- year-olds in Illinois fully vaccinated

https://dph.illinois.gov/covid19/vaccine/vaccine-data.html

As of 7/06/22


# COVID-19 Vaccines


Moderna

### Moderna

Age Group	Series How many shots, and when?	Dosage Compared to adult dose	Booster Are booster doses recommended?
MODERNA			
6 months-5 years	2 shots – 2 doses, 4 weeks apart	One-quarter	Not at this time
6-11 years	2 shots – 2 doses, 4 weeks apart	One-half	Not at this time
12-17 years	2 shots – 2 doses, 4 weeks apart	Equal	Not at this time

### Moderna






# COVID-19 Vaccines


Pfizer

### Pfizer

Age Group	Series How many shots, and when?	Dosage Compared to adult dose	Booster Are booster doses recommended?
PFIZER			
6 months-4 years	3 shots – 3 weeks between the first two doses, third dose at least 2 months after the second	One-tenth	Not at this time
5-11 years	2 shots – 2 doses, 3 weeks apart	One-third	Yes, at least 5 months after second shot
12-17 years	2 shots – 2 doses, 3 weeks apart	Equal	Yes, at least 5 months after second shot

### Pfizer Vials





### Most Common Side Effects

### ON THE ARM OR LEG EXPECT:

THROUGHOUT YOUR BODY:

•PAIN •REDNESS •SWELLING TIREDNESS
HEADACHE
MUSCLE PAIN
CHILLS
FEVER
NAUSEA



### V-Safe

A smartphone-based tool that uses text messaging and web surveys to give personalized health check-ins after patients receive a COVID-19 vaccine, found <u>here</u>

- ▷ Used to monitor vaccine safety
- Sends out vaccine reminders
- ▷ Does not give medical advice
- Cannot schedule COVID-19 vaccination appointments
- ▷ Not an official record of being vaccinated against COVID-19

Resources for people with disabilities

### CDC Website

- Videos
- Social Stories available in English and Spanish
  - How I get my <u>COVID-19 shot</u>
  - How I get my COVID-19 booster shot



### Resources

The Disability Information and Access Line (DIAL) connects callers to information about how to access the COVID-19 vaccine and related supports for people with disabilities.

DIAL connects callers to vaccine sites and provides information related to barriers to vaccination by referring callers to local and national disability resources.

Call 888-677-1199 from 8:00 AM to 7:00 pm or email <u>DIAL@n4a.org</u>. For more information, visit <u>acl.gov/dial</u>.



## Autistic Self Advocacy Network (ASAN) -<u>Videos</u> -<u>Fact Sheets in plain language</u>

## Thank You!